

# Fisheries Oceanography

November 2006 • Vol. 15 • Issue 6 • Page i-518

## ORIGINAL ARTICLES

- Retention mechanisms of white perch (*Morone americana*) and striped bass (*Morone saxatilis*) early-life stages in an estuarine turbidity maximum: an integrative fixed-location and mapping approach**  
E. W. NORTH, E. D. HOUDE .....429
- Hypoxia-based habitat compression of tropical pelagic fishes**  
ERIC D. PRINCE, C. PHILLIP GOODYEAR.....451
- Linking growth to environmental histories in central Baltic young-of-the-year sprat, *Sprattus sprattus*: an approach based on otolith microstructure analysis and hydrodynamic modelling**  
HANNES BAUMANN, HANS-HARALD HINRICHSSEN, RUDI VOSS, DANIEL STEPPUTTIS,  
WLODZIMIERZ GRYGIEL, LOTTE W. CLAUSEN, AXEL TEMMING .....465
- El Niño effects in the Palmyra Atoll region: oceanographic changes and bigeye tuna (*Thunnus obesus*) catch rate variability**  
EVAN A. HOWELL, DONALD R. KOBAYASHI .....477
- Temporal and spatial variability in growth and condition of dab (*Limanda limanda*) and sprat (*Sprattus sprattus*) larvae in the Irish Sea**  
ORLA LEE, BRET S. DANILOWICZ, MARK DICKEY-COLLAS .....490
- Distribution of jack mackerel (*Trachurus japonicus*) larvae and juveniles in the East China Sea, with special reference to the larval transport by the Kuroshio Current**  
CHIYUKI SASSA, YOSHINOBU KONISHI, KEN MORI .....508